

## Aberystwyth University

### *Morphological and physiological traits that explain yield response to drought stress in miscanthus*

Malinowska, Marta; Donnison, Iain; Robson, Paul

*Published in:*  
Agronomy

*DOI:*  
[10.3390/agronomy10081194](https://doi.org/10.3390/agronomy10081194)

*Publication date:*  
2020

*Citation for published version (APA):*

Malinowska, M., Donnison, I., & Robson, P. (2020). Morphological and physiological traits that explain yield response to drought stress in miscanthus. *Agronomy*, 10(8), [1194]. <https://doi.org/10.3390/agronomy10081194>

#### **Document License** CC BY

#### **General rights**

Copyright and moral rights for the publications made accessible in the Aberystwyth Research Portal (the Institutional Repository) are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Aberystwyth Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Aberystwyth Research Portal

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

tel: +44 1970 62 2400  
email: [is@aber.ac.uk](mailto:is@aber.ac.uk)

### Supplementary Table S1

Genotypes used in the experiment with information on species, country of origin and ploidy. *M. × giganteus* a commercial triploid hybrid between *M. sacchariflorus* and *M. sinensis*

WT code	species	country	ploidy
WAT03	<i>M. sacchariflorus</i>	Japan	4
WAT04	<i>M. sacchariflorus</i> (cv. Robustus)	China	2
WAT09	<i>M. × giganteus</i>	Japan	3
WAT10	<i>M. sinensis</i>	unkn.	unkn.
WAT11	<i>M. sinensis</i> (cv. Goliath)	Japan	3

### Supplementary Table S2

Drought resistance indices were calculated using the following relationships

$$SSI = 1 - (Y_s/Y_p) / 1 - (\bar{Y}_s/\bar{Y}_p)$$

$$STI = (Y_p + Y_s) / \bar{Y}_p^2$$


$$MP = (Y_p + Y_s) / 2$$

$$TOL = Y_p - Y_s$$

Where  $Y_s$  yield under stress,  $Y_p$  is the yield under irrigated condition  $\bar{Y}_s$  and  $\bar{Y}_p$  are the mean yields of all cultivars under stress and control conditions, respectively.

### Supplementary Figure S1

Principal component analysis (PCA) of morphology and biomass characteristics measured at harvests, of control (A) and plants under moderate drought stress (B; The overlapped variable names on the plot B are height, mprateH, and below-ground; The abbreviations used: WUE, water-use efficiency; below-ground, dry weight of below-ground biomass; above-ground, dry weight of above-ground biomass; height, the height of the main stem; noStems, averaged number of stem per plant; leafLength, the length of the youngest fully expanded leaf with ligule; leafWidth, the width of the youngest fully expanded leaf with ligule of the main stem; noLeaves, the number of leaves on the main stem; leafArea, area of the youngest fully leaf of the main stem; mprateLA, mean progression rate of the leaf area; mprateH, mean progression rate of the main stem.



### Supplementary Table S3

Analysis of variance on effects of drought stress (DS) treatment on the net transpiration rate (E), and net CO<sub>2</sub> assimilation (AN) of miscanthus genotypes, SE – measure of the uncertainty in the coefficient; t value is derived from the t-statistic and P-value indicates the statistical significance for a given term within the model. Significance of differences are shown as .P ≤ 0.1; \*P ≤ 0.05; \*\*P ≤ 0.01; \*\*\*P ≤ 0.001; ns = not significant.

Genotype	Source of variation	net CO <sub>2</sub> assimilation				net transpiration rate			
		Coefficient ± SE	t-val	P		Coefficient ± SE	t-val	P	
WAT03	(Intercept)	21,94 ± 2,98	7,34	ns		1,39 ± 0,26	5,18	ns	
	treatmentDS	4,88 ± 4,22	1,15	ns		0,61 ± 0,38	1,6	ns	
	TimePoint	-1,14 ± 1,42	-0,80	*		0,26 ± 0,12	2,04	ns	
	treatmentDS:TimePoint	-1,80 ± 2,01	-0,89	ns		-0,27 ± 0,18	-1,5	ns	
WAT04	(Intercept)	23,75 ± 4,17	5,68			1,34 ± 0,41	3,21	ns	
	treatmentDS	4,19 ± 5,86	0,71	**		0,70 ± 0,58	1,19	**	
	TimePoint	0,13 ± 1,74	0,08	.		0,41 ± 0,16	2,59	ns	
	treatmentDS:TimePoint	-4,59 ± 2,41	-1,90	.		-0,56 ± 0,22	-2,5	*	
WAT09	(Intercept)	27,32 ± 2,63	10,35	ns		1,83 ± 0,35	5,22	ns	
	treatmentDS	-0,80 ± 3,73	-0,21	***		-0,10 ± 0,49	-0,2	**	
	TimePoint	-1,64 ± 1,26	-1,29	**		0,22 ± 0,18	1,2	ns	
	treatmentDS:TimePoint	-2,18 ± 1,79	-1,22	ns		-0,22 ± 0,25	-0,9	ns	
WAT10	(Intercept)	34,13 ± 2,90	11,75	ns		2,92 ± 0,41	7,09	ns	
	treatmentDS	-5,47 ± 4,10	-1,33	ns		-0,78 ± 0,58	-1,3	ns	

WAT11	TimePoint	-7,10 ± 1,51	-4,68	***	-0,39 ± 0,18	-2,1	.
	treatmentDS:TimePoint	2,42 ± 2,14	1,12	ns	0,34 ± 0,26	1,29	ns
	(Intercept)	15,36 ± 3,09	4,95	ns	1,19 ± 0,24	4,91	ns
	treatmentDS	2,19 ± 4,29	0,51	.	0,14 ± 0,33	0,43	ns
	TimePoint	-1,58 ± 1,61	-0,98	**	0,05 ± 0,13	0,41	ns
	treatmentDS:TimePoint	-2,41 ± 2,19	-1,10		-0,18 ± 0,18	-1	

#### Supplementary Table S4

The mean values of water use efficiency and cumulative water use for each genotype under both treatments and the analysis of variance on the effect of drought stress on shoot water use efficiency (WUE), and the absolute water use compared to control; SEM – standard error of the mean; P-value indicates the statistical significance for a given term within the model. Significance of differences are shown as . $P \leq 0.1$ ; \* $P \leq 0.05$ ; \*\* $P \leq 0.01$ ; \*\*\* $P \leq 0.001$ ; ns = not significant..

	Water use efficiency (g kg <sup>-1</sup> )				Water use (kg)			
	Control	Drought	Treatment effect		Control	Drought	Treatment effect	
	Mean ±	Mean ±	Coefficient ± SE	P	Mean ±	Mean ±	Coefficient	P
WAT03	8.50 ±	12.59 ±	4.09 ± 1.03	**	9.92 ±	5.50 ±	-5.42 ± 0.76	***
WAT04	10.14 ±	16.93 ±	6.79 ± 0.74	***	6.47 ±	2.78 ±	-3.70 ± 0.86	**
WAT09	9.38 ±	13.42 ±	4.05 ± 0.71	***	8.70 ±	4.00 ±	-4.70 ± 1.02	**
WAT10	6.69 ±	15.57 ±	8.89 ± 0.92	***	7.60 ±	2.92 ±	-4.69 ± 0.75	***
WAT11	8.03 ±	17.32 ±	9.29 ± 0.65	***	7.17 ±	2.62 ±	-4.97 ± 1.42	***